

**QUASIPOTENTIAL IN THE FOURTH ORDER OF
PERTURBATION THEORY.
Unequal mass case.**

N.A. Boikova¹, V.V. Dvoeglazov¹, Yu.N. Tyukhtyaev ¹
R.N. Faustov²

Abstract

The quasipotential in the fourth order of perturbation theory is calculated in the Coulomb gauge for the unequal mass particles. It could be used for the future calculations of energy spectra in two-body systems.

Dubna, 1992

¹*Permanent address: Department of Theoretical and Nuclear Physics, Saratov State University and Scientific and Technical Center for Control and Use of Physical Fields and Radiation, Saratov 410071 RUSSIA*

²*Permanent address: Scientific Council "Cybernetics", Russian Academy of Sciences, Moscow 117333 RUSSIA*

At present the problem of an existence of the narrow resonances in (e^+e^-) - and (pp) -systems [1, 2] requires the adequate theoretical description [3]-[6]. The method proposed in the above papers is based on the quasipotential equation [7, 8] with the relativistic Coulomb potential modified with taking into account the binding effects in two-particle systems.

In this article the method of construction of the quasipotential by means of the two-time Green function $\hat{G}(x_a, x_b, t, y_a, y_b, t')$ is applied to the investigation of processes of the two-photon exchange.

The quasipotential

$$\hat{V} = F^{-1} - (\hat{G}^+)^{-1} \quad (1)$$

$$F = \hat{G}_0^+ = (2\pi)^3 \delta(\vec{p} - \vec{q}) (E - \sqrt{\vec{p}^2 + m_1^2} - \sqrt{\vec{p}^2 + m_2^2})^{-1} \quad (2)$$

was used earlier for the calculation of corrections to the Fermi energy of the hyperfine splitting and for the analysis of the fine structure energy in hydrogen-like atoms [9]-[11].

Recently the papers [12, 13]¹ devoted to the calculation of the $O(\alpha^6 \log \alpha)$ corrections in the two-body quantum electrodynamic systems have been appeared. The planned experiments will produce the question of eliminating the discrepancy in the results of calculation of the $O(\alpha^6 \log \alpha)$ corrections between [12] and [13].

The expressions for the quasipotential in the fourth order of perturbation theory (unequal mass case), which are presented in this paper, could be used for verifying the results [12, 13] and solving the problem. The indices C and T describe the successive exchange of the Coulomb photons and the transverse photons, respectively² The index "it" marks the expressions corresponding to the iteration diagrams. The index "x" marks the expressions corresponding to the cross diagrams (**Fig.1b,1e,1f,1h**).

$$\begin{aligned} \hat{V}_{CC}(\vec{p}, \vec{q}; E) = & -\frac{2\alpha^2}{\pi} u_1^*(\vec{p}) u_2^*(-\vec{p}) \int \frac{d\vec{k}}{k_p^2 k_q^2} \left\{ \frac{\Lambda_1^+(\vec{k}) \Lambda_2^+(-\vec{k})}{\epsilon_{1k} + \epsilon_{2k} - E} + \right. \\ & \left. + \frac{\Lambda_1^-(\vec{k}) \Lambda_2^-(-\vec{k})}{\epsilon_{1k} + \epsilon_{2k} + E} \right\} u_1(\vec{q}) u_2(-\vec{q}) \end{aligned} \quad (5)$$

$$\begin{aligned} \hat{V}_{CC}^\times(\vec{p}, \vec{q}; E) = & \frac{2\alpha^2}{\pi} u_1^*(\vec{p}) u_2^*(-\vec{p}) \int \frac{d\vec{k}}{k_p^2 k_q^2} \left\{ \frac{\Lambda_1^+(\vec{k}) \Lambda_2^-(\vec{k} - \vec{p} - \vec{q})}{\epsilon_{2p} + \epsilon_{1k} + \epsilon_{2kpq} + \epsilon_{2q} - E} + \right. \\ & \left. + \frac{\Lambda_1^-(\vec{k}) \Lambda_2^+(\vec{k} - \vec{p} - \vec{q})}{\epsilon_{1p} + \epsilon_{1k} + \epsilon_{2kpq} + \epsilon_{1q} - E} \right\} u_1(\vec{q}) u_2(-\vec{q}) \end{aligned} \quad (6)$$

¹The results of these papers are different for the positronium from each other:

$$\Delta E_{[12]} = \frac{1}{96} m \alpha^6 \log \alpha^{-1} (3 + 5\vec{\sigma}_1 \vec{\sigma}_2) \frac{\delta_{l0}}{n^3} \quad (3)$$

$$\Delta E_{[13]} = \frac{5}{96} m \alpha^6 \log \alpha^{-1} (3 + \vec{\sigma}_1 \vec{\sigma}_2) \frac{\delta_{l0}}{n^3} \quad (4)$$

² The Coulomb gauge is used.

$$\begin{aligned}
\hat{V}_{CT}(\vec{p}, \vec{q}; E) = & \frac{\alpha^2}{\pi} u_1^*(\vec{p}) u_2^*(-\vec{p}) \int \frac{d\vec{k}}{k_p^2 k_q} \left\{ \frac{\Lambda_1^+(\vec{k}) \Lambda_2^+(-\vec{k})}{\epsilon_{1k} + \epsilon_{2k} - E} \left(\frac{1}{k_q + \epsilon_{2k} + \epsilon_{1q} - E} + \right. \right. \\
& + \left. \frac{1}{k_q + \epsilon_{1k} + \epsilon_{2q} - E} \right) + \frac{\Lambda_1^-(\vec{k}) \Lambda_2^-(-\vec{k})}{\epsilon_{1k} + \epsilon_{2k} + E} \left(\frac{1}{k_q + \epsilon_{1k} + \epsilon_{1q}} + \frac{1}{k_q + \epsilon_{2k} + \epsilon_{2q}} \right) - \\
& - \frac{\Lambda_1^-(\vec{k}) \Lambda_2^+(-\vec{k})}{(k_q + \epsilon_{2k} + \epsilon_{1q} - E)(k_q + \epsilon_{1k} + \epsilon_{1q})} - \frac{\Lambda_1^+(\vec{k}) \Lambda_2^-(-\vec{k})}{(k_q + \epsilon_{1k} + \epsilon_{2q} - E)(k_q + \epsilon_{2k} + \epsilon_{2q})} \Big\} \times \\
& \times \Gamma_{12}(\vec{k} - \vec{q}) u_1(\vec{q}) u_2(-\vec{q})
\end{aligned} \tag{7}$$

$$\begin{aligned}
\hat{V}_{CT}^\times(\vec{p}, \vec{q}; E) = & \frac{\alpha^2}{\pi} u_1^*(\vec{p}) u_2^*(-\vec{p}) \int \frac{d\vec{k}}{k_p^2 k_q} \left\{ \frac{\Lambda_1^+(\vec{k}) \Gamma_{12}(\vec{k} - \vec{q}) \Lambda_2^+(\vec{k} - \vec{p} - \vec{q})}{(k_q + \epsilon_{1p} + \epsilon_{2kpq} - E)(k_q + \epsilon_{1k} + \epsilon_{2q} - E)} + \right. \\
& + \frac{\Lambda_1^-(\vec{k}) \Gamma_{12}(\vec{k} - \vec{q}) \Lambda_2^-(\vec{k} - \vec{p} - \vec{q})}{(k_q + \epsilon_{2p} + \epsilon_{2kpq})(k_q + \epsilon_{1k} + \epsilon_{1q})} - \frac{\Lambda_1^+(\vec{k}) \Gamma_{12}(\vec{k} - \vec{q}) \Lambda_2^-(\vec{k} - \vec{p} - \vec{q})}{\epsilon_{2p} + \epsilon_{1k} + \epsilon_{2kpq} + \epsilon_{2q} - E} \times \\
& \times \left(\frac{1}{k_q + \epsilon_{2p} + \epsilon_{2kpq}} + \frac{1}{k_q + \epsilon_{1k} + \epsilon_{2q} - E} \right) - \frac{\Lambda_1^-(\vec{k}) \Gamma_{12}(\vec{k} - \vec{q}) \Lambda_2^+(\vec{k} - \vec{p} - \vec{q})}{\epsilon_{1p} + \epsilon_{1k} + \epsilon_{2kpq} + \epsilon_{1q} - E} \times \\
& \times \left. \left(\frac{1}{k_q + \epsilon_{1p} + \epsilon_{2kpq} - E} + \frac{1}{k_q + \epsilon_{1k} + \epsilon_{1q}} \right) \right\} u_1(\vec{q}) u_2(-\vec{q})
\end{aligned} \tag{8}$$

$$\begin{aligned}
\hat{V}_{TT}(\vec{p}, \vec{q}; E) = & -\frac{\alpha^2}{2\pi} u_1^*(\vec{p}) u_2^*(-\vec{p}) \int \frac{d\vec{k}}{k_p k_q} \Gamma_{12}(\vec{p} - \vec{k}) \left\{ \frac{\Lambda_1^+(\vec{k}) \Lambda_2^+(-\vec{k})}{\epsilon_{1k} + \epsilon_{2k} - E} \times \right. \\
& \times \left[\frac{1}{k_p + k_q + \epsilon_{1p} + \epsilon_{2q} - E} \left(\frac{1}{k_p + \epsilon_{1p} + \epsilon_{2k} - E} + \frac{1}{k_q + \epsilon_{1k} + \epsilon_{2q} - E} \right) + \right. \\
& + \frac{1}{k_p + k_q + \epsilon_{2p} + \epsilon_{1q} - E} \left(\frac{1}{k_p + \epsilon_{2p} + \epsilon_{1k} - E} + \frac{1}{k_q + \epsilon_{2k} + \epsilon_{1q} - E} \right) + \\
& + \left. \frac{1}{(k_p + \epsilon_{2p} + \epsilon_{1k} - E)(k_q + \epsilon_{1k} + \epsilon_{2q} - E)} + \frac{1}{(k_p + \epsilon_{1p} + \epsilon_{2k} - E)(k_q + \epsilon_{2k} + \epsilon_{1q} - E)} \right] + \\
& + \frac{\Lambda_1^-(\vec{k}) \Lambda_2^-(-\vec{k})}{\epsilon_{1k} + \epsilon_{2k} + E} \left[\frac{1}{k_p + k_q + \epsilon_{1p} + \epsilon_{2q} - E} \left(\frac{1}{k_p + \epsilon_{1p} + \epsilon_{2k} - E_1 + E_2} + \right. \right. \\
& + \left. \frac{1}{k_q + \epsilon_{1k} + \epsilon_{2q} + E_1 - E_2} \right) + \frac{1}{k_p + k_q + \epsilon_{2p} + \epsilon_{1q} - E} \times \\
& \times \left(\frac{1}{k_p + \epsilon_{2p} + \epsilon_{1k} + E_1 - E_2} + \frac{1}{k_q + \epsilon_{2k} + \epsilon_{1q} - E_1 + E_2} \right) + \\
& + \left. \frac{1}{(k_p + \epsilon_{2p} + \epsilon_{1k} + E_1 - E_2)(k_q + \epsilon_{1k} + \epsilon_{2q} + E_1 - E_2)} + \right. \\
& + \left. \frac{1}{(k_p + \epsilon_{1p} + \epsilon_{2k} - E)(k_q + \epsilon_{2k} + \epsilon_{1q} - E)} \right] + \frac{\Lambda_1^-(\vec{k}) \Lambda_2^+(-\vec{k})}{-\epsilon_{1k} + \epsilon_{2k} - E} \times \\
& \times \left[\frac{1}{k_p + k_q + \epsilon_{1p} + \epsilon_{2q} - E} \left(\frac{1}{k_p + \epsilon_{1p} + \epsilon_{2k} - E} - \frac{1}{k_p + \epsilon_{1p} + \epsilon_{1k}} \right) + \right.
\end{aligned}$$

$$\begin{aligned}
& + \frac{1}{k_p + k_q + \epsilon_{2p} + \epsilon_{1q} - E} \left(\frac{1}{k_q + \epsilon_{2k} + \epsilon_{1q} - E} - \frac{1}{k_q + \epsilon_{1k} + \epsilon_{1q}} \right) + \\
& + \frac{1}{(k_p + \epsilon_{1p} + \epsilon_{2k} - E)(k_q + \epsilon_{2k} + \epsilon_{1q} - E)} - \frac{1}{(k_p + \epsilon_{1p} + \epsilon_{1k})(k_q + \epsilon_{1k} + \epsilon_{1q})} \Big] + \\
& + \frac{\Lambda_1^+(\vec{k})\Lambda_2^-(\vec{k})}{\epsilon_{1k} - \epsilon_{2k} - E} \left[\frac{1}{k_p + k_q + \epsilon_{1p} + \epsilon_{2q} - E} \left(\frac{1}{k_q + \epsilon_{1k} + \epsilon_{2q} - E} - \frac{1}{k_q + \epsilon_{2k} + \epsilon_{2q}} \right) + \right. \\
& + \frac{1}{k_p + k_q + \epsilon_{2p} + \epsilon_{1q} - E} \left(\frac{1}{k_p + \epsilon_{2p} + \epsilon_{1k} - E} - \frac{1}{k_p + \epsilon_{2p} + \epsilon_{2k}} \right) + \\
& + \left. \frac{1}{(k_p + \epsilon_{2p} + \epsilon_{1k} - E)(k_q + \epsilon_{1k} + \epsilon_{2q} - E)} - \frac{1}{(k_p + \epsilon_{2p} + \epsilon_{2k})(k_q + \epsilon_{2k} + \epsilon_{2q})} \right] \Big\} \times \\
& \times \Gamma_{12}(\vec{k} - \vec{q})u_1(\vec{q})u_2(-\vec{q})
\end{aligned} \tag{9}$$

$$\begin{aligned}
\hat{V}_{TT}^\times(\vec{p}, \vec{q}; E) = & -\frac{\alpha^2}{2\pi} u_1^*(\vec{p})u_2^*(-\vec{p}) \int \frac{d\vec{k}}{k_p k_q} \alpha_{1i}(\delta_{ij} - \frac{(\vec{p} - \vec{k})_i(\vec{p} - \vec{k})_j}{(\vec{p} - \vec{k})^2}) \times \\
& \times \left\{ \Lambda_1^+(\vec{k})\Gamma_{12}(\vec{k} - \vec{q})\Lambda_2^+(\vec{k} - \vec{p} - \vec{q}) \left[\frac{1}{k_p + k_q + \epsilon_{1k} + \epsilon_{2kpq} - E} \left(\frac{1}{k_p + \epsilon_{2p} + \epsilon_{1k} - E} + \right. \right. \right. \\
& + \frac{1}{k_q + \epsilon_{1p} + \epsilon_{2kpq} - E} \left(\frac{1}{k_p + \epsilon_{2kpq} + \epsilon_{1q} - E} + \frac{1}{k_q + \epsilon_{1k} + \epsilon_{2q} - E} \right) + \\
& + \frac{1}{(k_p + k_q + \epsilon_{1p} + \epsilon_{2q} - E)(k_q + \epsilon_{1p} + \epsilon_{2kpq} - E)(k_q + \epsilon_{1k} + \epsilon_{2q} - E)} + \\
& + \left. \frac{1}{(k_p + k_q + \epsilon_{2p} + \epsilon_{1q} - E)(k_p + \epsilon_{2p} + \epsilon_{1k} - E)(k_p + \epsilon_{2kpq} + \epsilon_{1q} - E)} \right] + \\
& + \Lambda_1^-(\vec{k})\Gamma_{12}(\vec{k} - \vec{q})\Lambda_2^-(\vec{k} - \vec{p} - \vec{q}) \left[\frac{1}{k_p + k_q + \epsilon_{2kpq} + \epsilon_{1k} + E} \left(\frac{1}{k_p + \epsilon_{1p} + \epsilon_{1k}} + \right. \right. \\
& + \frac{1}{k_q + \epsilon_{2p} + \epsilon_{2kpq}} \left(\frac{1}{k_q + \epsilon_{1k} + \epsilon_{1q}} + \frac{1}{k_p + \epsilon_{2kpq} + \epsilon_{2q}} \right) + \\
& + \frac{1}{(k_p + k_q + \epsilon_{1p} + \epsilon_{2q} - E)(k_p + \epsilon_{1p} + \epsilon_{1k})(k_p + \epsilon_{2kpq} + \epsilon_{2q})} + \\
& + \left. \frac{1}{(k_p + k_q + \epsilon_{2p} + \epsilon_{1q} - E)(k_q + \epsilon_{2p} + \epsilon_{2kpq})(k_q + \epsilon_{1k} + \epsilon_{1q})} \right] - \\
& - \Lambda_1^+(\vec{k})\Gamma_{12}(\vec{k} - \vec{q})\Lambda_2^-(\vec{k} - \vec{p} - \vec{q}) \left[\frac{1}{\epsilon_{2p} + \epsilon_{1k} + \epsilon_{2kpq} + \epsilon_{2q} - E} \left(\frac{1}{k_p + \epsilon_{2kpq} + \epsilon_{2q}} + \right. \right. \\
& + \frac{1}{k_p + \epsilon_{2p} + \epsilon_{1k} - E} \left(\frac{1}{k_q + \epsilon_{2p} + \epsilon_{2kpq}} + \frac{1}{k_q + \epsilon_{1k} + \epsilon_{2q} - E} \right) + \\
& + \frac{1}{(k_p + k_q + \epsilon_{1p} + \epsilon_{2q} - E)(k_p + \epsilon_{2kpq} + \epsilon_{2q})(k_q + \epsilon_{1k} + \epsilon_{2q} - E)} + \\
& + \left. \frac{1}{(k_p + k_q + \epsilon_{2p} + \epsilon_{1q} - E)(k_p + \epsilon_{2p} + \epsilon_{1k} - E)(k_q + \epsilon_{2p} + \epsilon_{2kpq})} \right] - \\
& - \Lambda_1^-(\vec{k})\Gamma_{12}(\vec{k} - \vec{q})\Lambda_2^+(\vec{k} - \vec{p} - \vec{q}) \left[\frac{1}{\epsilon_{1p} + \epsilon_{1k} + \epsilon_{2kpq} + \epsilon_{1q} - E} \left(\frac{1}{k_p + \epsilon_{1p} + \epsilon_{1k}} + \right. \right. \\
& + \frac{1}{k_p + \epsilon_{2kpq} + \epsilon_{1q} - E} \left(\frac{1}{k_p + \epsilon_{1k} + \epsilon_{1q}} + \frac{1}{k_q + \epsilon_{1p} + \epsilon_{2kpq} - E} \right) +
\end{aligned}$$

$$\begin{aligned}
& + \frac{1}{(k_p + k_q + \epsilon_{1p} + \epsilon_{2q} - E)(k_p + \epsilon_{1p} + \epsilon_{1k})(k_q + \epsilon_{1p} + \epsilon_{2kpq} - E)} + \\
& + \frac{1}{(k_p + k_q + \epsilon_{2p} + \epsilon_{1q} - E)(k_p + \epsilon_{2kpq} + \epsilon_{1q} - E)(k_q + \epsilon_{1k} + \epsilon_{1q})} \Big] \Big\} \times \\
& \times \alpha_{2j} u_1(\vec{q}) u_2(-\vec{q})
\end{aligned} \tag{10}$$

$$\hat{V}_{CC}^{it}(\vec{p}, \vec{q}; E) = -\frac{2\alpha^2}{\pi} u_1^*(\vec{p}) u_2^*(-\vec{p}) \int \frac{d\vec{k}}{k_p^2 k_q^2} \frac{\Lambda_1^+(\vec{k}) \Lambda_2^+(-\vec{k})}{\epsilon_{1k} + \epsilon_{2k} - E} u_1(\vec{q}) u_2(-\vec{q}) \tag{11}$$

$$\begin{aligned}
\hat{V}_{CT}^{it}(\vec{p}, \vec{q}; E) &= \frac{\alpha^2}{\pi} u_1^*(\vec{p}) u_2^*(-\vec{p}) \int \frac{d\vec{k}}{k_p^2 k_q} \frac{\Lambda_1^+(\vec{k}) \Lambda_2^+(-\vec{k})}{\epsilon_{1k} + \epsilon_{2k} - E} \times \\
&\times \left(\frac{1}{k_q + \epsilon_{1k} + \epsilon_{2q} - E} + \frac{1}{k_q + \epsilon_{2k} + \epsilon_{1q} - E} \right) \Gamma_{12}(\vec{k} - \vec{q}) u_1(\vec{q}) u_2(-\vec{q})
\end{aligned} \tag{12}$$

$$\begin{aligned}
\hat{V}_{TT}^{it}(\vec{p}, \vec{q}; E) &= -\frac{\alpha^2}{2\pi} u_1^*(\vec{p}) u_2^*(-\vec{p}) \int \frac{d\vec{k}}{k_p k_q} \Gamma_{12}(\vec{p} - \vec{k}) \frac{\Lambda_1^+(\vec{k}) \Lambda_2^+(-\vec{k})}{\epsilon_{1k} + \epsilon_{2k} - E} \times \\
&\times \left(\frac{1}{k_p + \epsilon_{2p} + \epsilon_{1k} - E} + \frac{1}{k_p + \epsilon_{1p} + \epsilon_{2k} - E} \right) \left(\frac{1}{k_q + \epsilon_{1k} + \epsilon_{2q} - E} + \right. \\
&+ \left. \frac{1}{k_q + \epsilon_{2k} + \epsilon_{1q} - E} \right) \Gamma_{12}(\vec{k} - \vec{q}) u_1(\vec{q}) u_2(-\vec{q})
\end{aligned} \tag{13}$$

Here $\Lambda^\pm$ are the projecting operators, $k_p = |\vec{p} - \vec{k}|$, $k_q = |\vec{k} - \vec{q}|$, $k_{pq} = |\vec{k} - \vec{p} - \vec{q}|$, $\Gamma_{12}(\vec{k}) = \vec{\alpha}_1 \vec{\alpha}_2 - (\vec{\alpha}_1 \vec{k})(\vec{\alpha}_2 \vec{k})/\vec{k}^2$, $\vec{\alpha}_i = \gamma_i^0 \vec{\gamma}_i$ and $\gamma_\mu = (\gamma^0, \vec{\gamma})$ – Dirac's matrices.

The expressions corresponding to **Fig.1d,1e,1k** are obtained from the expressions (7), (8), (12) by means of the substitutions $p \leftrightarrow q$ and the displacement of the projecting operators and Γ_{12} . The quasipotentials for the diagrams with a vertex part and a self-energy part are easily deduced from the one-photon exchange quasipotential presented e.g in [6, 9, 11, 14, 15].

We are grateful to B. A. Arbuzov, E. E. Boos, V. G. Kadyshevsky and V. I. Savrin for useful discussions and to N. B. Skachkov for the hospitality.

References

- [1] Troyan Yu. A. et al. JINR Communications Nos.R1-90-52, 78, 79, 139, Dubna: JINR, 1990;
- [2] Cowan T. , Howell R. H., Rohatgi R. R., Fajans J. "Nucl. Instr. and Meth.", 1991, v. B56/57, part I, p. 599-603 (XI Int. Conf. on the application of accelerators in research and industry, Denton, TX (USA), 5-8 Nov. 1990);
- [3] Arbuzov B. A. et al. "Pisma JETP", 1989, v.50, No.5, p.236-238;
- [4] Arbuzov B. A. et al. "Teor. Mat. Fiz", 1990, v. 83, No. 2, p. 175-185;
- [5] Arbuzov B.A. et al. "Mod. Phys. Lett.", 1990, v. A5, No. 18, p. 1441-1449;

- [6] Arbuzov B. A. et al. IHEP preprint 90-124, Protvino: IHEP, 1990;
- [7] Logunov A. N., Tavkhelidze A. N. "Nuovo Cim.", 1963, v. 29, No. 2, p. 380-399;
- [8] Nguyen Van Hieu, Faustov R. N. "Nucl. Phys.", 1964, v. 53, No. 2, p. 337-344;
- [9] Levchenko N. A., Tyukhtyaev Yu. N., Faustov R. N. JINR Communications No. R2-12355, Dubna: JINR, 1979 ; "Yad. Fiz.", 1980, v. 32, No. 6(12), p. 1656-1662;
- [10] Boikova N. A., Tyukhtyaev Yu. N., Faustov R. N. JINR Communications No. R2-81-582, Dubna: JINR, 1981;
- [11] Boikova N. A. et al. "Teor. Mat. Fiz.", 1991, v.89, No.2, p. 228-237;
- [12] Fell R. N. "Phys. Rev. Lett.", 1992, v.68, No.1 ,p.25 - 28;
- [13] Khriplovich I. B., Milstein A. I., Yelkhovsky A. S. CERN preprint CERN-TH 6420/92, Geneva: CERN, 1992.
- [14] Arkhipov A. A. "Teor. Mat. Fiz", 1990, v.83, No.2, p.247-267; ibid, No.3, p. 358-373;
- [15] Kapshaj V. N., Savrin V. I., Skachkov N. B. "Teor. Mat. Fiz.", 1986,v.69, No.3, p.400-410; Dey E. A., Kapshaj V. N., Tyumenkov G. Yu. "Acta Phys. Pol.", 1990, v. B21, No. 6, p. 449 - 456.




Fig. 1. The diagrams for calculation of the quasipotential in the fourth order of perturbation theory(unequal mass case). The dot line corresponds to the Coulomb photon; the dashed line, to the transverse photon.